

Datasheet

Cryopan I

Serum-free Freezing Medium

Product	Description	Catalogue-No.	Size
Cryopan I	Serum-free freezing medium, with 10 % DMSO	P07-92010	10 ml
		P07-92050	50 ml
		P07-92100	100 ml
		P07-92500	500 ml

Product description

Cryopan I is a serum-free, ready-to-use freezing medium for animal and human cells (adherent and suspension cells). It is especially suitable for cells from serum-free culture.

Storage conditions

Storage conditions: - 20°C
 Stability: 2 years from date of production
 Filling: 10 ml, 50 ml, 100 ml, 500 ml, other sizes on request

Composition


Cryopan I consists of an optimized mixture of salts, sugar, 10 % DMSO and additional antifreeze-substances.

Suitability

Cryopan I is for the cryoconservation of human and animal cells.

Special Advantages

Due to its serum-free formulation, Cryopan is especially suitable for the conservation of serum-free cultured cells. The optimized composition guarantees a high cell viability after the thawing process. Best results are attained for all cell types including primary human cells and human cell lines.



Instructions for Use

1. Freezing cells with Cryopan I

For optimal results only vital cells in the log-growth phase should be used.

- Thaw Cryopan I and store till using at 2-8°C.
- Trypsinate adherent cells, transfer the cells into the culture medium, stop the trypsin activity with trypsin inhibitor and centrifuge.
- Discard the supernatant and wash the cell pellet in PBS (without Ca²⁺/Mg²⁺).
- After an additional centrifugation step (100 – 200 g, 5 - 10 minutes) transfer the cells into PBS and determine cell number and cell vitality by trypan blue cell viability staining.
- Centrifuge the cells again and discard the supernatant.
- Transfer the cells into the cold Cryopan I (5x10⁵ - 2x10⁶ cells/ml Cryopan I).
- Suspend the cells carefully mixing the suspension by repeated pipetting until there are no more cell clumps.
- Refill the cell suspension into labelled cryotubes (0,5 - 1,5ml/tube).
- Freeze the cells in an automatically or manually controlled cryo freezer. The optimal freezing rate is approximately 1°C/minute.
- Alternatively, put the tubes for 15 minutes into a refrigerator, so that the freezing medium can penetrate into the cells. After this step freeze the tubes at -20°C for 2 hours box and put them into the vapour phases of liquid nitrogen over night.
- Store cryotubes in a cryotank with liquid nitrogen.

2. Thawing cells

- Remove the cryotubes from the cryotank and thaw them as soon as possible in warm water (< 1 minute).
- Disinfect the exterior of the cryotubes with alcohol and convict the cells under sterile conditions into a centrifuge tube with 10 ml growth medium and mix it carefully.
- Centrifuge the cells (150 - 200 g, 5 - 10 minutes).
- Discard the supernatant and recover the cells into the designated culture medium. Determine the cell viability by an appropriate method, like FACS or trypan blue cell viability staining.

Technical support

For technical support, questions or remarks please contact your local PAN-Biotech partner or the technical department of PAN-Biotech via email (info@pan-biotech.com) or phone +49-8543-601630.

FOR RESEARCH USE ONLY! Not approved for human or animal diagnostic or therapeutic procedures.